




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K	A2	(11) International Publication Number: WO 98/18429
		(43) International Publication Date: 7 May 1998 (07.05.98)
<p>(21) International Application Number: PCT/EP97/05774</p> <p>(22) International Filing Date: 18 October 1997 (18.10.97)</p> <p>(30) Priority Data: 08/739,840 30 October 1996 (30.10.96) US</p> <p>(71) Applicant: SCHERING AKTIENGESELLSCHAFT [DE/DE]; D-13342 Berlin (DE).</p> <p>(72) Inventors: KAUSER, Katalin; 1299 Raton Court, El Sobrante, CA 94803 (US). RUBANYI, Gabor; 2426 Heritage Oaks Drive, Alamo, CA 94507 (US).</p>		<p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>

(54) Title: A METHOD FOR LOWERING PLASMA LEVELS OF LIPOPROTEIN(A)

(57) Abstract

A method of lowering plasma levels of Lp(a) or inhibiting the generation of apo(a) in a human by administering an effective amount of a compound of formula (I) wherein: R¹ is hydrogen or alkyl; A, B, D, E, G, R² and R³ each are hydrogen, and R⁴ is hydrogen in the β -position; or R² with R⁴, R⁴ with A, A with R³, B with D, D with E, or G with R² is a bond, and the other substituents are each hydrogen; or R² is alkyl in the β -position and A, B, D, E, G, R³ and R⁴ are each hydrogen; or R³ is alkoxy in the β -position and A, B, D, E, G, R² and R⁴ are each hydrogen; or R⁴ is hydrogen in the α -position and A, B, D, E, G, R² and R³ are each hydrogen; and R⁵ is methyl or ethyl, is disclosed.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A METHOD FOR LOWERING PLASMA LEVELS OF LIPOPROTEIN(A)

Field of the Invention

This invention is directed to a method for lowering plasma levels of lipoprotein(a) in humans. In particular, this invention is directed to the use of certain 17-difluoromethylene-estratrienes to lower plasma levels of lipoprotein(a) in humans, thereby being useful in the treatment and prevention of premature occlusive arterial disease.

BACKGROUND OF THE INVENTION

Lipoprotein(a) (Lp(a)) is an unique lipoprotein found in the plasma of only human, some sub-human primates and the European hedgehog. Lp(a) is considered an independent risk factor for premature occlusive arterial diseases, such as atherosclerosis, coronary artery disease, myocardial infarction, cerebral infarction (stroke), peripheral artery disease and restenosis following balloon angioplasty. The structure of Lp(a) is closely related to LDL in that it consists of LDL with an additional disulfide-linked apolipoprotein, apo(a). Apo(a) is covalently bound to the glycoprotein Apo B100, an integral part of LDL. Apo B100 allows the LDL molecule to carry the hydrophobic cholesterol in the plasma and tissue fluids. In contrast, apo(a) is water soluble and does not bind lipid. The major site of synthesis of plasma apo(a) appears to be the liver. However, it is currently unknown where Lp(a) is assembled. LDL and apo(a) may be secreted independently of each other, and the assembly may take place in the plasma. Such a mechanism could explain why only a fraction of LDL, which varies from subject to subject, is complexed with apo(a). This fraction is determined by the amount of apo(a) secreted rather than by the LDL concentration in plasma. From *in vivo* turnover studies of Lp(a) in humans, it has been concluded that differences in plasma Lp(a) concentrations among individuals are a result of differences in synthesis rather than differences in catabolism of Lp(a).

Elevated plasma levels of Lp(a) is associated with premature occlusive arterial disease, such as coronary heart disease (CHD) and stroke. In addition, the association between concentrations of Lp(a) in serum and coronary artery disease (CAD) has been shown. Furthermore, the level of Lp(a) in the plasma has been shown to independently correlate with the rate of restenosis (after balloon angioplasty), peripheral arterial occlusive disease, diabetic vasculopathies in Type II diabetes mellitus, increased CHD risk in

patients undergoing continuous ambulatory peritoneal dialysis, and stenosis of bypass grafts.

Surprisingly, the associations of Lp(a) with other known risk factors for CHD, for example, plasma lipid (triglycerides, LDL, HDL) levels, dietary fat intake, hypertension, diabetes, overweight and cigarette smoking, have been consistently weak or absent and do not appear to explain the relation of Lp(a) to CHD.


It is widely believed that the level of Lp(a) in the plasma is under strong genetic control, with non-genetic factors, including gender, diabetes mellitus and end-stage renal disease, having some influence. Lp(a) levels in men and women differ significantly and are influenced by physiological conditions such as puberty and menopause when production of sex hormones is altered. Lp(a) levels in females continue to rise with age, particularly in the premenopausal years, while in males they plateau at about 25 years of age. In addition, Lp(a) plasma levels have been found to be higher in postmenopausal than premenopausal women of comparable age. A gender related difference can also be observed in studies which have correlated the levels of Lp(a) with the extent of angiographically documented CAD. Lp(a) plasma levels have also been shown to be more predictable of CAD in females than in males.

The use of steroid hormones (e.g., estrogens, progesterone and testosterone) to lower plasma levels of Lp(a) has been shown. For example, Lp(a) plasma concentrations have been shown to be influenced by the administration of anabolic steroids, progesterone, estrogen, and combination of estrogen and progestogen to postmenopausal women. In males affected by prostatic cancer, estrogen therapy lowered Lp(a) plasma levels by 50%, while orchidectomy treatment slightly increased the plasma concentration of Lp(a), suggesting that estrogen and testosterone exert a regulatory role on serum Lp(a) levels in men.

We have now discovered that certain 17-difluoromethylene-estratrienes are effective in inhibiting the generation of apo(a) and are, consequently, effective in lowering plasma levels of Lp(a), and are therefore useful in treating disease-states which are alleviated by inhibiting the generation of apo(a) or by lowering the plasma levels of Lp(a). These compounds have not been disclosed previously as having this ability.

SUMMARY OF THE INVENTION

Accordingly, one aspect of this invention is directed to a method of lowering plasma levels of Lp(a) in a human, which method comprises administering to a human in need thereof an effective amount of a compound of formula (I):



wherein:

R¹ is hydrogen or alkyl;

A, B, D, E, G, R² and R³ each are hydrogen, and R⁴ is hydrogen in the β -position; or

10 R² with R⁴, R⁴ with A, A with R³, B with D, D with E, or G with R² is a bond, and the other substituents are each hydrogen; or

R² is alkyl in the β -position and A, B, D, E, G, R³ and R⁴ are each hydrogen; or

R³ is alkoxy in the β -position and A, B, D, E, G, R² and R⁴ are each hydrogen; or

R⁴ is hydrogen in the α -position and A, B, D, E, G, R² and R³ are each hydrogen; and

15 R⁵ is methyl or ethyl.

A further aspect of this invention is directed to a method of inhibiting the generation of apo(a) in a human, which method comprises administering a human in need thereof an effective amount of a compound of formula (I) as defined above.

20 Another aspect of this invention is directed to a method of treating a human having a disease-state which is alleviated by reducing the plasma levels of Lp(a), which method comprises administering to the human in need thereof a therapeutically effective amount of a compound of formula (I) as defined above.

A further aspect of this invention is directed to a method of treating a human having a disease-state which is alleviated by inhibiting the generation of apo(a), which method

comprises administering to the human in need thereof a therapeutically effective amount of a compound of formula (I) as defined above.

Another aspect of this invention is directed to a pharmaceutical composition useful in lowering plasma levels of Lp(a) in a human, which composition comprises an effective amount of a compound of formula (I) as defined above and a pharmaceutically acceptable excipient.

A further aspect of this invention is directed to a pharmaceutical composition useful in inhibiting the generation of apo(a) in a human, which composition comprises an effective amount of a compound of formula (I) as defined above and a pharmaceutically acceptable excipient.

Another aspect of this invention is directed to a method of inhibiting the generation of apo(a) *in vitro* or *in vivo* by administering a compound of formula (I) as defined above.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used in the specification and appended claims, unless specified to the contrary, the following terms have the meaning indicated:

"Alkyl" refers to a straight or branched chain monovalent radical consisting solely of carbon and hydrogen atoms, containing no unsaturation and having from one to ten carbon atoms, e.g., methyl, ethyl, propyl, butyl, 1-methylethyl (*iso*-propyl), 1,1-dimethylethyl (*tert*-butyl), pentyl, *n*-hexyl, *n*-decyl, and the like.

"Alkoxy" refers to a radical of the formula -OR_a where R_a is alkyl as defined above, e.g., methoxy, ethoxy, *n*-propoxy, *n*-heptoxy, and the like.

"Premature occlusive arterial disease" refers to those disease-states, for the purposes of this invention, which are characterized by the progressive narrowing and eventual closure of an artery or the process of progressive narrowing and eventual closing an artery, and which a) develop earlier in life than normal and b) are associated with high levels of Lp(a) in the plasma; and includes, *inter alia*, arteriosclerosis, atherosclerosis, coronary artery disease, peripheral artery disease, myocardial infarction, stroke, restenosis and bypass graft stenosis.

"Effective amount" refers to that amount of a compound of formula (I) which, when administered to a human, is sufficient to lower the plasma level of Lp(a) or to inhibit the generation of apo(a).

"Therapeutically effective amount" refers to that amount of a compound of formula (I) which, when administered to a human in need thereof, is sufficient to effect treatment, as defined below, for disease-states alleviated by the reduction of plasma levels of Lp(a) or by the inhibition of the generation of apo(a). The amount of a compound of formula (I) which constitutes a "therapeutically effective amount" will vary depending on the compound, the disease-state and its severity, and the age of the human to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

"Treating" or "treatment" as used herein cover the treatment of a disease-state in a human, which disease-state is alleviated by the reduction of plasma levels of Lp(a) or the inhibition of the generation of apo(a); and include:

- (i) preventing the disease-state from occurring in a human, in particular, when such human is predisposed to the disease-state but has not yet been diagnosed as having it;
- (ii) inhibiting the disease-state, *i.e.*, arresting its development; or
- (iii) relieving the disease-state, *i.e.*, causing regression of the disease-state.


Brief Description of the Drawings

Various other objects, features and attendant advantages of the invention will be more fully appreciated as the same becomes better understood when considered in conjunction with the accompanying drawings wherein:

FIGURE 1 shows plasma apo(a) levels and uterine dry weight of ovariectomized female apo(a) transgenic mice after treatment with either a compound of formula (I) or vehicle.

Preferred Embodiments

Of the various aspects of this invention as set forth above, a preferred aspect is the method of lowering plasma levels of Lp(a) in a human, which method comprises administering to a human in need thereof an effective amount of a compound of formula (I) of the following formula:



namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

Another preferred aspect of this invention is the method of inhibiting the generation of apo(a) in a human, which method comprises administering to a human in need thereof an effective amount of the compound of formula (I) as described above.

Another preferred aspect of this invention is the method of treating a human having a premature occlusive vascular disease which is alleviated by the reduction of plasma levels of Lp(a), which method comprises administering to the human in need thereof a therapeutically effective amount of the compound of formula (I) as described above.

Another preferred aspect of this invention is the method of treating a human having a premature occlusive vascular disease which is alleviated by the inhibition of the generation of apo(a), which method comprises administering to the human in need thereof a therapeutically effective amount of the compound of formula (I) as described above.

Utility of the Invention

This invention is directed to methods for lowering the plasma level of Lp(a) or for inhibiting the generation of apo(a) in humans, particularly in women without significant side effect on the endometrium. As discussed above, there is a direct correlation between high levels of Lp(a) and premature occlusive arterial disease. However, the exact physiological function of either apo(a) or Lp(a) is still being investigated. Apparently, even the near absence of Lp(a) from plasma does not cause a deficiency syndrome or any kind of disease. Also, Lp(a) and apo(a) are apparently absent in the plasma from most species evolutionarily below Old World monkeys. However, various pathological functions have recently been postulated for both Lp(a) and apo(a) based on the results of certain *in vitro* and *in vivo* studies as indicators of premature occlusive arterial diseases.

For example, it is known that from the total pool of cholesterol that is transported in the blood, only a very small fraction is associated with Lp(a). This small pool, however, is believed to act as "quick response" reserve for areas where rapid cell turnover, active membrane biosynthesis or inflammatory processes occur, such as the atheromatous plaque. Indeed, high levels of Lp(a) have been found to be localized in the vascular intima (significantly more than the circulating plasma level would indicate), especially in the intima at sites of atheromatous lesions and stenotic saphenous vein bypass grafts.

Another proposed function involves the endothelium, which is normally antithrombogenic. The luminal surface of endothelial cells binds plasminogen, which is activated by tissue plasminogen activator (TPA) on the endothelial cell surface to produce the antithrombogenic agent, plasmin. It is postulated that Lp(a) binds competitively to the plasminogen binding site and reduces the amount of plasmin generated by TPA. This can lead to reduced antithrombogenic and antifibrinolytic properties of the endothelium.

It has also been shown that Lp(a) binds to fibrin, which may prevent degradation of an existing thrombus by plasminogen and may contribute to Lp(a) accumulation in atherosclerotic plaques.

It has also been shown that Lp(a) prolongs the time required for fibrinolysis in an assay where fibrinolytic activity is stimulated by a fibrinolytic agent, e.g., TPA/streptokinase. The mechanism for this appears to be an inhibition of the conversion of plasminogen to plasmin by the fibrinolytic agent. Further evidence for its potential role in atherosclerosis is provided by studies showing monocyte adhesion glycoprotein expression by endothelial cells as a result of treatment with Lp(a). In addition, a recent study showed that Lp(a) may contribute to activation of immune response in atheromatous lesions.

Recently, it has been shown that transgenic mice expressing the human apo(a) gene develop atherosclerotic lesions in response to a high fat diet, which suggests a causative link between atherogenesis and apo(a), a necessary component of Lp(a). See, Lawn, R.M. *et al.*, *Nature* 1992, Vol. 360, pp. 670-672.


To demonstrate the utility of the compounds of the invention as therapeutic agents for treating disease-states which are alleviated by the reduction of plasma Lp(a) levels, we evaluated the compounds for their ability to inhibit the generation of apo(a) in mice made transgenic for the human apo(a) gene (see Frazer, K.A. *et al.*, *Nature Genetics* (1995), Vol. 9, pp. 424-431, for description of the transgenic mice). In particular, the transgenic mice were treated with the compounds of the invention and their plasma levels of apo(a)

were determined. In addition, the uterine dry weight of each mouse was determined to demonstrate any side effect of the treatment on the uterus.

As illustrated in **FIGURE 1**, a significant decrease in the plasma apo(a) levels of the treated mice was observed. In addition, as illustrated in **FIGURE 1**, the compounds appear to have little or no significant effect on the uterus, as evidenced by the uterine dry weight of the mice after treatment. Since apo(a) is a necessary component of Lp(a), a reduction in the level of apo(a) would also reduce the level of Lp(a) in the plasma. Compounds which reduce plasma levels of Lp(a) would, upon administration, alleviate those disease-states which are characterized by high levels of Lp(a), such as premature occlusive arterial diseases.

B. Preparation of the Compounds of the Invention

The compounds used in this invention are selected from the following formula (I):



wherein:

R¹ is hydrogen or alkyl;

A, B, D, E, G, R² and R³ each are hydrogen, and R⁴ is hydrogen in the β-position; or R² with R⁴, R⁴ with A, A with R³, B with D, D with E, or G with R² is a bond, and the other substituents are each hydrogen; or


R² is alkyl in the β-position and A, B, D, E, G, R³ and R⁴ are each hydrogen; or

R³ is alkoxy in the β-position and A, B, D, E, G, R² and R⁴ are each hydrogen; or

R⁴ is hydrogen in the α-position and A, B, D, E, G, R² and R³ are each hydrogen; and

R⁵ is methyl or ethyl.

These compounds are prepared treating the corresponding 17-oxo compound of the following formula:



wherein R^1 is an hydroxy protecting group and A, B, D, E, G, R^2 , R^3 , R^4 and R^5 are the same as described above in the Summary of the Invention, with difluoromethyl-diphenylphosphine oxide or diethyl(difluoromethyl)phosphonate in the presence of a strong base in
 5 a aprotic solvent at a reflux temperature of from about 50°C to about 100°C, and then optionally the 3-hydroxy protective group is cleaved under the action of an acid to produce the corresponding 3-hydroxy group, which can be optionally etherified.

The 3-hydroxy protective group is either a radical that can easily be cleaved in acid or basic environment, such as, for example, a tetrahydropyranyl (THP) group or a silyl
 10 group that is substituted with three identical, two identical, or three different straight or branched chain alkyl and/or aryl (such as phenyl or naphthyl) radicals, such as, for example, the trimethyl, *t*-butyldimethyl, methyldiphenyl, or *t*-butyldiphenylsilyl group or a methyl group, which can be removed, however, only under more drastic conditions.

Lithium diisopropylamide, sodium hydride, potassium *t*-butoxide, butyllithium and the
 15 like are suitable as a strong base.

The reaction of the 17-oxo compound is carried out in an aprotic solvent, such as, for example, tetrahydrofuran, dimethyl sulfoxide, dimethylformamide, dioxane, or a mixture thereof.

The conditions for the cleavage of the 3-hydroxy protective group depends on its
 20 nature. Protective groups such as tetrahydropyranyl or a silyl radical can be removed under the action of a weak acid such as oxalic acid or an acidic ion exchanger, while the methyl group can be cleaved under the action of strong Lewis acids, e.g., dibutylaluminum hydride.

The etherification of the free 3-hydroxy group is carried out with a reagent that
 25 yields R^1 groups in a manner known to those skilled in the art.

The production of the 3-hydroxy-protected-17-oxo compounds used in the preparation of the compounds is carried out by reaction of the corresponding known 3-hydroxy compound with dihydropyran under the effect of *para*-toluenesulfonic acid in tetrahydrofuran or other methods known to those skilled in the art for protection of hydroxy groups.

Detailed synthesis of these compounds is described in detail in PCT Published Patent Application WO 96/28462 (Bohlmann *et al.*), which is incorporated in full by reference herein.

10 C. General Administration

Administration of the compounds of the invention, in pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration or agents for serving similar utilities. Thus, administration can be, for example, orally, nasally, parenterally, topically, transdermally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages. The compositions will include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, *etc.*

Generally, depending on the intended mode of administration, the pharmaceutically acceptable compositions will contain about 1% to about 99% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and 99% to 1% by weight of a suitable pharmaceutical excipient. Preferably, the composition will be about 5% to 75% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, with the rest being suitable pharmaceutical excipients.

The preferred route of administration is oral, using a convenient daily dosage regimen which can be adjusted according to the degree of severity of the disease-state to be treated. For such oral administration, a pharmaceutically acceptable composition containing a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, is formed by the incorporation of any of the normally employed, pharmaceutically acceptable excipients, such as, for example, pharmaceutical grades of mannitol, lactose, starch, pregelatinized starch, magnesium stearate, sodium saccharine, talcum, cellulose ether

derivatives, glucose, gelatin, sucrose, citrate, propyl gallate, and the like. Such compositions take the form of solutions, suspensions, tablets, pills, capsules, powders, sustained release formulations and the like.

Preferably such compositions will take the form of capsule, caplet or tablet and
5 therefore will also contain a diluent such as lactose, sucrose, dicalcium phosphate, and the like; a disintegrant such as croscarmellose sodium or derivatives thereof; a lubricant such as magnesium stearate and the like; and a binder such as a starch, gum acacia, polyvinylpyrrolidone, gelatin, cellulose ether derivatives, and the like.

The compounds of the invention, or their pharmaceutically acceptable salts, may
10 also be formulated into a suppository using, for example, about 0.5% to about 50% active ingredient disposed in a carrier that slowly dissolves within the body, e.g., polyoxyethylene glycols and polyethylene glycols (PEG), e.g., PEG 1000 (96%) and PEG 4000 (4%).

Liquid pharmaceutically administrable compositions can, for example, be prepared
by dissolving, dispersing, etc., a compound(s) of the invention (about 0.5% to about 20%),
15 or a pharmaceutically acceptable salt thereof, and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol and the like, to thereby form a solution or suspension.

If desired, a pharmaceutical composition of the invention may also contain minor
amounts of auxiliary substances such as wetting or emulsifying agents, Ph buffering
20 agents, antioxidants, and the like, such as, for example, citric acid, sorbitan monolaurate, triethanolamine oleate, butylated hydroxytoluene, etc.

Actual methods of preparing such dosage forms are known, or will be apparent, to
those skilled in this art; for example, see *Remington's Pharmaceutical Sciences*, 18th Ed.,
(Mack Publishing Company, Easton, Pennsylvania, 1990). The composition to be
25 administered will, in any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state alleviated by the reduction of plasma levels of Lp(a) or by the inhibition of the generation of apo(a) in accordance with the teachings of this invention.

The compounds of the invention, or their pharmaceutically acceptable salts, are
30 administered in a therapeutically effective amount which will vary depending upon a variety of factors including the activity of the specific compound employed; the metabolic stability and length of action of the compound; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug

combination; the severity of the particular disease-states; and the patient undergoing treatment. Generally, a therapeutically effective daily dose is from about 0.14 mg to about 14.3 mg/kg of body weight per day of a compound of the invention, or a pharmaceutically acceptable salt thereof; preferably, from about 0.7 mg to about 10 mg/kg of body weight per day; and most preferably, from about 1.4 mg to about 7.2 mg/kg of body weight per day. For example, for administration to a 70 kg person, the dosage range would be from about 10 mg to about 1.0 gram per day of a compound of the invention, or a pharmaceutically acceptable salt thereof, preferably from about 50 mg to about 700 mg per day, and most preferably from about 100 mg to about 500 mg per day.

The following specific examples are provided as a guide to assist in the practice of the invention, and are not intended as a limitation on the scope of the invention.

EXAMPLE 1

Effect of a Compound of Formula (I) in Apo(a) Transgenic Mice

A. Ovariectomized female apo(a) transgenic mice were randomized into three groups, each group containing 3 animals. The test compound was 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol. The mice were treated with daily subcutaneous injections of 1 mg/kg of the test compound in vehicle, 10 mg/kg of the test compound in vehicle, and vehicle (benzylbenzoate:castor oil, 1:9). Blood samples were collected *via* tail bleeding at the end of the first week and *via* heart puncture at the end of the second week of treatment when the mice were sacrificed. At sacrifice, the uteri of the animals were dissected, cleaned from adherent connective tissues, placed on aluminum foil and dried in an oven for 24 hours. At the time dry organ weight was determined. Apo(a) levels were determined from plasma using the Macra ELISA Kit (Stretegic Diagnostic).

Results of the study, as illustrated in **FIGURE 1**, demonstrated that the test compound significantly lowered apo(a) levels at 1 mg/kg and 10 mg/kg dose, whereas significant uterotrophic activity ($*p < 0.05$) was only measured at the 10 mg/kg dose. This dissociation in the efficacy of the test compound to lower apo(a) levels and increase uterine weight demonstrates the test compound feasibility as an apo(a)/Lp(a) lowering agent without significant side effects on the uterus.

EXAMPLE 2

This example illustrates the preparation of representative pharmaceutical compositions for oral administration containing a compound used in the invention, e.g., 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol:

5	A.	<u>Ingredients</u>	<u>% wt./wt.</u>
		Compound of the invention	20.0%
		Lactose	79.5%
		Magnesium stearate	0.5%

The above ingredients are mixed and dispensed into hard-shell gelatin capsules

10 containing 100 mg each.

	B.	<u>Ingredients</u>	<u>% wt./wt.</u>
		Compound of the invention	20.0%
		Magnesium stearate	0.9%
		Starch	8.6%
15		Lactose	69.6%
		PVP (polyvinylpyrrolidone)	0.9%

The above ingredients with the exception of the magnesium stearate are combined and granulated using water as a granulating liquid. The formulation is then dried, mixed with the magnesium stearate and formed into tablets with an appropriate tableting

20 machine.

	C.	<u>Ingredients</u>	
		Compound of the invention	0.1 g
		Propylene glycol	20.0 g
		Polyethylene glycol 400	20.0 g
25		Polysorbate 80	1.0 g
		Water	q.s. 100 mL

The compound of the invention is dissolved in propylene glycol, polyethylene glycol 400 and polysorbate 80. A sufficient quantity of water is then added with stirring to provide 100 mL of the solution which is filtered and bottled.

30	D.	<u>Ingredients</u>	<u>% wt./wt.</u>
		Compound of the invention	20.0%
		Peanut Oil	78.0%
		Span 60	2.0%

The above ingredients are melted, mixed and filled into soft elastic capsules.

E.	<u>Ingredients</u>	<u>% wt./wt.</u>
	Compound of the invention	1.0%
	Methyl or carboxymethyl cellulose	2.0%
5	0.9% saline	q.s. 100 mL

The compound of the invention is dissolved in the cellulose/saline solution, filtered and bottled for use.

EXAMPLE 3

- 10 This Example illustrates the preparation of a representative pharmaceutical composition for parenteral administration containing a compound used in the invention, e.g., 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol:

	<u>Ingredients</u>	
	Compound of the invention	0.02 g
15	Propylene glycol	20.0 g
	Polyethylene glycol 400	20.0 g
	Polysorbate 80	1.0 g
	0.9% Saline solution	q.s. 100 mL

- 20 The compound of the invention is dissolved in propylene glycol, polyethylene glycol 400 and polysorbate 80. A sufficient quantity of 0.9% saline solution is then added with stirring to provide 100 mL of the I.V. solution which is filtered through a 0.2 m membrane filter and packaged under sterile conditions.

EXAMPLE 4

- 25 This Example illustrates the preparation of a representative pharmaceutical composition in suppository form containing a compound used in the invention, e.g., 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol:

	<u>Ingredients</u>	<u>% wt./wt.</u>
	Compound of the invention	1.0%
30	Polyethylene glycol 1000	74.5%
	Polyethylene glycol 4000	24.5%

The ingredients are melted together and mixed on a steam bath, and poured into molds containing 2.5 g total weight.

EXAMPLE 5

This Example illustrates the preparation of a representative pharmaceutical composition for insufflation containing a compound used in the invention, e.g.,

5 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol:

<u>Ingredients</u>	<u>% wt./wt.</u>
Micronized compound of the invention	1.0%
Micronized lactose	99.0%

The ingredients are milled, mixed, and packaged in an insufflator equipped with a
10 dosing pump.

EXAMPLE 6

This Example illustrates the preparation of a representative pharmaceutical composition in nebulized form containing a compound used in the invention, e.g.,

15 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol:

<u>Ingredients</u>	<u>% wt./wt.</u>
Compound of the invention	0.005%
Water	89.995%
Ethanol	10.000%

20 The compound of the invention is dissolved in ethanol and blended with water. The formulation is then packaged in a nebulizer equipped with a dosing pump.

EXAMPLE 7

25 This Example illustrates the preparation of a representative pharmaceutical composition in aerosol form containing a compound used in the invention, e.g.,

17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol:


<u>Ingredients</u>	<u>% wt./wt.</u>
Compound of the invention	0.10%
Propellant 11/12	98.90%
30 Oleic acid	1.00%

The compound of the invention is dispersed in oleic acid and the propellants. The resulting mixture is then poured into an aerosol container fitted with a metering valve.

While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a
5 particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

WHAT IS CLAIMED IS:

1. A method of lowering plasma levels of Lp(a) in a human, which method comprises administering to the human in need thereof an effective amount of a compound of formula (I):



wherein:

R¹ is hydrogen or alkyl;

A, B, D, E, G, R² and R³ each are hydrogen, and R⁴ is hydrogen in the β -position; or R² with R⁴, R⁴ with A, A with R³, B with D, D with E, or G with R² is a bond, and the other substituents are each hydrogen; or


R² is alkyl in the β -position and A, B, D, E, G, R³ and R⁴ are each hydrogen; or

R³ is alkoxy in the β -position and A, B, D, E, G, R² and R⁴ are each hydrogen; or

R⁴ is hydrogen in the α -position and A, B, D, E, G, R² and R³ are each hydrogen; and


R⁵ is methyl or ethyl.

2. The method of Claim 1 which comprises administering a compound of the formula:



namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

3. A method of inhibiting the generation of apo(a) in a human, which method comprises administering a human in need thereof an effective amount of a compound of formula (I):



wherein:

R^1 is hydrogen or alkyl;

A, B, D, E, G, R^2 and R^3 each are hydrogen, and R^4 is hydrogen in the β -position; or R^2 with R^4 , R^4 with A, A with R^3 , B with D, D with E, or G with R^2 is a bond, and the other substituents are each hydrogen; or


R^2 is alkyl in the β -position and A, B, D, E, G, R^3 and R^4 are each hydrogen; or

R^3 is alkoxy in the β -position and A, B, D, E, G, R^2 and R^4 are each hydrogen; or

R^4 is hydrogen in the α -position and A, B, D, E, G, R^2 and R^3 are each hydrogen; and


R^5 is methyl or ethyl.

4. The method of Claim 3 which comprises administering a compound of the formula:



namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

5. A method of treating a human having a disease-state which is alleviated by the reduction of plasma levels of Lp(a), which method comprises administering to the human in need thereof a therapeutically effective amount of a compound of formula (I):



wherein:

R¹ is hydrogen or alkyl;


A, B, D, E, G, R² and R³ each are hydrogen, and R⁴ is hydrogen in the β -position; or

R² with R⁴, R⁴ with A, A with R³, B with D, D with E, or G with R² is a bond, and the other substituents are each hydrogen; or

R² is alkyl in the β -position and A, B, D, E, G, R³ and R⁴ are each hydrogen; or


R^3 is alkoxy in the β -position and A, B, D, E, G, R^2 and R^4 are each hydrogen; or
 R^4 is hydrogen in the α -position and A, B, D, E, G, R^2 and R^3 are each hydrogen; and
 R^5 is methyl or ethyl.

6. The method of Claim 5 which comprises administering a compound of the formula:




namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

7. The method of Claim 5 wherein the disease-state is premature occlusive vascular disease.
8. The method of Claim 7 wherein the disease-state is stroke.
9. The method of Claim 7 wherein the disease-state is coronary heart disease.
10. The method of Claim 7 wherein the disease-state is diabetes mellitus.
11. The method of Claim 7 wherein the disease-state is atherosclerosis.
12. The method of Claim 7 which comprises administering a compound of the formula:



namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

13. A method of treating a human having a disease-state which is alleviated by the inhibition of generation of apo(a), which method comprises administering to the human in need thereof a therapeutically effective amount of a compound of formula (I):



wherein:

R^1 is hydrogen or alkyl;

A, B, D, E, G, R^2 and R^3 each are hydrogen, and R^4 is hydrogen in the β -position; or R^2 with R^4 , R^4 with A, A with R^3 , B with D, D with E, or G with R^2 is a bond, and the other substituents are each hydrogen; or


R^2 is alkyl in the β -position and A, B, D, E, G, R^3 and R^4 are each hydrogen; or

R^3 is alkoxy in the β -position and A, B, D, E, G, R^2 and R^4 are each hydrogen; or

R^4 is hydrogen in the α -position and A, B, D, E, G, R^2 and R^3 are each hydrogen; and


R^5 is methyl or ethyl.

14. The method of Claim 13 which comprises administering a compound of the formula:




namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

15. The method of Claim 13 wherein the disease-state is premature occlusive vascular disease.
16. The method of Claim 15 wherein the disease-state is stroke.
17. The method of Claim 15 wherein the disease-state is coronary heart disease.
18. The method of Claim 15 wherein the disease-state is diabetes mellitus.
19. The method of Claim 15 wherein the disease-state is atherosclerosis.
20. The method of Claim 15 which comprises administering a compound of the formula:



namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

21. A pharmaceutical composition useful in treating a human having a disease-state which is alleviated by the reduction of plasma levels of Lp(a), which composition comprises a therapeutically effective amount of a compound of formula (I):



wherein:

R^1 is hydrogen or alkyl;

A, B, D, E, G, R^2 and R^3 each are hydrogen, and R^4 is hydrogen in the β -position; or R^2 with R^4 , R^4 with A, A with R^3 , B with D, D with E, or G with R^2 is a bond, and the other substituents are each hydrogen; or


R^2 is alkyl in the β -position and A, B, D, E, G, R^3 and R^4 are each hydrogen; or

R^3 is alkoxy in the β -position and A, B, D, E, G, R^2 and R^4 are each hydrogen; or

R^4 is hydrogen in the α -position and A, B, D, E, G, R^2 and R^3 are each hydrogen; and


R^5 is methyl or ethyl.

22. The composition of Claim 21 which comprises a therapeutically effective amount of a compound of the formula:



namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

23. A pharmaceutical composition useful in treating a human having a disease-state which is alleviated by the inhibition of the generation of apo(a), which composition comprises a therapeutically effective amount of a compound of formula (I):



wherein:


R¹ is hydrogen or alkyl;

A, B, D, E, G, R² and R³ each are hydrogen, and R⁴ is hydrogen in the β -position; or
R² with R⁴, R⁴ with A, A with R³, B with D, D with E, or G with R² is a bond, and the other
substituents are each hydrogen; or

R² is alkyl in the β -position and A, B, D, E, G, R³ and R⁴ are each hydrogen; or


R^3 is alkoxy in the β -position and A, B, D, E, G, R^2 and R^4 are each hydrogen; or
 R^4 is hydrogen in the α -position and A, B, D, E, G, R^2 and R^3 are each hydrogen; and
 R^5 is methyl or ethyl.

24. The composition of Claim 23 which comprises a therapeutically effective amount of a compound of the formula:



namely, 17-difluoromethylene-18-methylestra-1,3,5(10)-trien-3-ol.

1/1

**FIGURE 1**

Plasma levels of apo(a) in treated and control mice after 2-week treatment.

Values represent means \pm SEM of plasma apo(a) levels (left Y axis) and uterine weight (right Y axis) measured in 3 mice in each group.

Asterisk (*) indicates statistically significant ($p < 0.05$) difference between the treated and control group.




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/56		A3	(11) International Publication Number: WO 98/18429
			(43) International Publication Date: 7 May 1998 (07.05.98)
(21) International Application Number: PCT/EP97/05774		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 18 October 1997 (18.10.97)			
(30) Priority Data: 08/739,840 30 October 1996 (30.10.96) US			
(71) Applicant: SCHERING AKTIENGESELLSCHAFT [DE/DE]; D-13342 Berlin (DE).		Published With international search report.	
(72) Inventors: KAUSER, Katalin; 1299 Raton Court, El Sobrante, CA 94803 (US). RUBANYI, Gabor; 2426 Heritage Oaks Drive, Alamo, CA 94507 (US).		(88) Date of publication of the international search report: 8 October 1998 (08.10.98)	

(54) Title: A METHOD FOR LOWERING PLASMA LEVELS OF LIPOPROTEIN(A)

(57) Abstract

A method of lowering plasma levels of Lp(a) or inhibiting the generation of apo(a) in a human by administering an effective amount of a compound of formula (I) wherein: R¹ is hydrogen or alkyl; A, B, D, E, G, R² and R³ each are hydrogen, and R⁴ is hydrogen in the β -position; or R² with R⁴, R⁴ with A, A with R³, B with D, D with E, or G with R² is a bond, and the other substituents are each hydrogen; or R² is alkyl in the β -position and A, B, D, E, G, R³ and R⁴ are each hydrogen; or R³ is alkoxy in the β -position and A, B, D, E, G, R² and R⁴ are each hydrogen; or R⁴ is hydrogen in the α -position and A, B, D, E, G, R² and R³ are each hydrogen; and R⁵ is methyl or ethyl, is disclosed.



(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/05774

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/56

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 28462 A (SCHERING AG ; BOHLMANN ROLF (DE); RUBANYL GABOR (US)) 19 September 1996 cited in the application see abstract see page 6, paragraph 3 - page 7, last paragraph; claims; examples ---	1-24
A	R.M. LAWN ET AL.: "ATHEROGENESIS IN TRANSGENIC MICE EXPRESSING HUMAN APOLIPOPROTEIN(a)" NATURE, vol. 360, 1992, pages 670-672, XP002070686 cited in the application see the whole document ---	1-24

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

7 July 1998

Date of mailing of the international search report

31.07.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoff, P

1

PCT/EP 97/05774

1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 97/05774

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-20
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-20 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/05774

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9628462 A	19-09-1996	DE 19509729 A	19-09-1996
		AU 5107896 A	02-10-1996
		CA 2215201 A	19-09-1996
		CZ 9702810 A	18-02-1998
		EP 0815120 A	07-01-1998
		NO 974206 A	13-11-1997
		PL 322199 A	19-01-1998
		SI 9620030 A	28-02-1998
EP 0320438 A	14-06-1989	SK 121697 A	04-02-1998
		DE 3741800 A	15-06-1989
		AU 2665288 A	08-06-1989
		CA 1324374 A	16-11-1993
		DE 3874865 A	29-10-1992
		ES 2045176 T	16-01-1994
		GR 3006579 T	30-06-1993
		JP 1193295 A	03-08-1989
		JP 2636913 B	06-08-1997
		US 5124321 A	23-06-1992